

The synchrotron foreground and CMB temperature–polarization cross correlation power spectrum from the first year WMAP data

G. Bernardi,¹ E. Carretti,¹ R. Fabbri,² C. Sbarra,¹ S. Cortiglioni¹

¹*INAF-IASF Bologna, Via Gobetti 101, I-40129 Bologna, Italy*

²*Dipartimento di Fisica, Università di Firenze, Via Sansone 1, I-50019 Sesto Fiorentino (FI), Italy*

25 August 2005

ABSTRACT

We analyse the temperature-polarization cross-correlation in the Galactic synchrotron template that we recently developed, and between the template and CMB temperature maps derived from WMAP data. Since the polarized synchrotron template itself uses WMAP data, we can estimate residual synchrotron contamination in the CMB C_ℓ^{TE} angular spectrum. While C_2^{TE} appears to be contaminated by synchrotron, no evidence for contamination is found in the multipole range which is most relevant for the fit of the cosmological optical depth.

Key words: cosmic microwave background, polarization, (cosmology:) diffuse radiation, method: data analysis

1 INTRODUCTION

The WMAP experiment measuring the temperature-polarization cross-correlation power spectrum of the Cosmic Microwave Background (CMB) found an excess of power at large angular scales ($\ell < 10$), which has been interpreted as evidence for an early reionization (Kogut et al. 2003). A clean measurement of the cosmological signal relies on a successful removal of the foregrounds, which on large angular scales are mainly generated by dust, free-free and synchrotron emissions from the Galaxy. In particular, the synchrotron radiation is the main polarized foreground at WMAP frequencies. According to Bennett et al. (2003) the CMB maps used to compute the angular power spectrum C_ℓ^T have negligible foreground contamination, thanks to the wide frequency coverage of the WMAP experiment and a safe foreground subtraction achieved with fits of foreground templates. Also Kogut et al. (2003) claimed that the contamination in the Q, V and W bands is low when the Galactic plane is cut out and the C_ℓ^{TE} power spectrum of the CMB is free of foreground contamination.

However, several groups have performed independent analyses of the WMAP data to address the foreground contamination on the CMB maps. Tegmark, de Oliveira-Costa & Hamilton (2003, hereafter TOH) claimed to have obtained a CMB map^{*} cleaner than the one of the WMAP team. Naselsky et al. (2003) applied a phase analysis to the

internal linear combination map obtained by the WMAP team, showing some residual foreground contamination. Also, Naselsky et al. (2004) compared the analysis of the internal linear combination map obtained by the WMAP team’s analysis with TOH’s, and found evidence for a residual contamination in the low-multipole power spectrum region. Dineen & Coles (2004a, 2004b) used the cross-correlation between the rotation measures of extragalactic radio sources and the CMB maps to identify a possible foreground residual; they found evidence for that in both the WMAP and TOH CMB maps. However, these works cannot tell us whether the foreground residual may affect the C_ℓ^{TE} power spectrum to a significant extent, and it looks harder to improve the cross-correlation analysis of Kogut et al. (2003) because polarization maps have not been provided yet by WMAP. The issue of possible foreground contamination on C_ℓ^{TE} is however very important in the light of the reported anomalies in WMAP’s large-scale output, including North-South asymmetries (Eriksen et al. 2004, Hansen et al. 2004b, Land and Maguejo 2005) and multipole alignments (TOH, de Oliveira-Costa et al. 2004, Copi et al. 2003). As far as we are concerned about the robustness of the C_ℓ^{TE} power spectrum and the inferences on cosmological reionization, the most troublesome result is due to Hansen et al. (2004a), according to which the high optical depth ascribed to the cosmological medium by WMAP’s team should originate from the Southern (Galactic or Ecliptic) hemisphere.

In the light of the above results, it is necessary to investigate further the possible impact of foregrounds on the

* <http://www.hep.upenn.edu/~max/>




Figure 1. The synchrotron C_ℓ^{TE} power spectrum at 60 GHz (solid line) compared to the CMB C_ℓ^{TE} from Kogut et al. (2003) (asterisks). Also reported are the $C_\ell^{TCMBE SYNCH}$ power spectra for the WMAP CMB map (dashed line) and for the TOH CMB map (dash-dotted).

C_ℓ^{TE} power spectrum. Due to the absence of measured polarization maps at microwave frequencies, uncertainties on the foreground contamination in polarization are greater than in total intensity. In this Letter we want to tackle the problem of the foreground contamination on the CMB C_ℓ^{TE} power spectrum on large angular scales using the templates of synchrotron polarized emission developed by Bernardi et al. (2003) and Bernardi et al. (2004, hereafter B04). The B04 template in particular is expected to be much more accurate than the earlier one which relies on the extrapolation of surveys at ~ 1 GHz. However, its superiority derives from the use of WMAP’s total intensity maps.

2 SYNCHROTRON AND CMB C_ℓ^{TE} POWER SPECTRA

B04 provided Q and U template maps of the Galactic synchrotron emission at 23 GHz, which is obtained from the 23 GHz total intensity synchrotron map released by the WMAP team (by using the polarization direction field of starlight as well as a polarization horizon model). B04 also derived a Galactic map of the synchrotron spectral index, which is used to scale the polarization template to higher frequencies. In the present work, the same spectral index map is used to scale the WMAP 23-GHz total intensity synchrotron map released by the WMAP team. We choose a frequency of 60 GHz, which can be regarded as an approximate mean of the frequencies of the WMAP QVW data set. Synchrotron T , Q and U maps (with T the antenna temperature) are generated at an angular resolution of 7° due to the limitation of the B04 template. We are then able to compute the 60-GHz synchrotron C_ℓ^{TE} power spectrum by integration of the two-point correlation function. This procedure (see, for the implementation, Sbarra et al. 2003) allows to properly account for the incomplete sky coverage of the B04 template, and for the kp2 Galactic-Plane mask applied to the synchrotron maps (for details, see Bennett et al. 2003). A

similar procedure allows us to investigate cross-correlations between the CMB and the synchrotron template.

Figure 1 shows the synchrotron C_ℓ^{TE} power spectrum at 60 GHz and the CMB C_ℓ^{TE} power spectrum measured by WMAP. The error bars on the synchrotron spectrum only account for a variation $\Delta\alpha = \pm 0.2$ of the frequency spectral index of the synchrotron emission. Therefore they represent the uncertainty on the overall normalization of the fiducial synchrotron template at 60 GHz, and do not account for statistical errors. Clearly, the 60-GHz synchrotron C_ℓ^{TE} is much smaller than the corresponding CMB spectrum in the range $\ell = 3 - 10$ where most information about cosmological reionization is encoded. The situation of course would be still better at 90 GHz. We note that the synchrotron quadrupole C_2^{TE} has a large and negative value and this indicates a potential source of contamination for the CMB quadrupole; however, this very fact does not indicate any inadequacy in WMAP team’s technique of foreground removal. On the other hand, reasonably strong evidence for a residual contamination can be provided by a cross-correlation between the CMB temperature and the synchrotron polarization fields, both being derived from WMAP’s data.

In order to cross-correlate our Q and U templates with CMB anisotropy, we use two different CMB maps. The first one is obtained by averaging the Q , V and W maps released by the WMAP team after foreground subtraction (we refer to this as the WMAP CMB map); the other one is the CMB map produced by TOH. The $C_\ell^{TCMBE SYNCH}$ power spectra computed for both these maps are also shown in Fig. 1. Both power spectra show very similar behaviours for $\ell > 2$. We find no evidence of a CMB-synchrotron correlation in the range $\ell = 4 - 10$. The multipole $\ell = 3$ shows a cross-correlation which at 60 GHz is comparable to the CMB C_3^{TE} . This should not be so disturbing after all, since the large reionization optical depth is essentially generated by slightly larger ℓ ’s. The most intriguing feature is still the behaviour of the quadrupole. When the synchrotron template is correlated with the WMAP CMB map we find a large ($\sim 8 \mu\text{K}^2$) negative value, whereas the use of the TOH CMB map leads to a relatively lower ($\sim 5 \mu\text{K}^2$) but positive value. For comparison, the CMB quadrupole is $C_2^{TE} \sim 3 \mu\text{K}^2$, a factor 2–4 lower than the magnitude of $C_2^{TCMBE SYNCH}$.

The discrepancy between cross-correlation quadrupoles is not surprising, since TOH already noted that their temperature quadrupole C_2^T is significantly different from the one found by the WMAP team. This discrepancy, as well as the overall behaviour of the cross-correlation power spectra, is better understood by inspection of the CMB-synchrotron cross-correlation function. The latter is defined by

$$C^{TQ}(\theta) = \sum_{ij} I_i Q_j^r, \quad (1)$$

where the Stokes parameter Q^r is computed in the frame of the great circle connecting pixels (i, j) . Figure 2 shows the cross-correlation functions derived from WMAP and TOH maps and their difference. It is interesting to note that the cross-correlation functions have the same behaviour up to $\theta \sim 20^\circ$ but they differ significantly for larger angular scales. There is a clear evidence for strongly correlated signals between the WMAP CMB map and the polarized synchrotron at angular scales $\theta > 50^\circ$. The absolute maximum of $C^{TQ}(\theta)$ is $\sim 2 \mu\text{K}^2$ for $\theta \sim 110^\circ$. The

[tbp]




Figure 2. Top panel: The correlation function between the synchrotron Q^r and the CMB temperature maps provided by WMAP’s team (solid line) and by TOH (dashed line). Bottom panel: The difference between the two correlation functions.

cross-correlation function derived from the TOH CMB map is somewhat weaker, and it has an opposite sign at several scales, especially the scales that are the most relevant in the determination of $C_2^{TCMBESYNCH}$. However, the difference between the two correlation functions approximately has a very simple angle dependence $\Delta C^{TQ}(\theta) \propto \sin\theta$ and is as large as $\sim 2\mu\text{K}^2$ for $\theta \sim 90^\circ$. This fully accounts for the discrepancy found for $C_2^{TCMBESYNCH}$. The rather moderate decrease of both $C^{TQ}(\theta)$ and $C_2^{TCMBESYNCH}$ achieved by using the TOH temperature map seems to imply that a residual contamination by synchrotron emission survives on the largest angular scales even in the TOH treatment.

3 CONCLUSIONS

The main purpose of this *Letter* is the study of the possible synchrotron contamination of the CMB C_ℓ^{TE} power spectrum derived from the first year WMAP release. The comparison between the CMB and the synchrotron template C_ℓ^{TE} power spectra shows that the contamination in the 60-GHz CMB C_ℓ^{TE} should be negligible for $3 \leq \ell \leq 10$. The inspection of the cross-correlation spectrum $C_\ell^{TCMBESYNCH}$ (as well as the analysis of the cross-correlation function) reinforces this conclusion at least in the range $4 \leq \ell \leq 10$. On the other hand, the high values of the synchrotron quadrupole C_2^{TE} and of the CMB-synchrotron cross-correlated signal on large angular scales ($\theta > 50^\circ$) suggest a residual synchrotron contamination. We emphasize that the B04 template uses the WMAP total intensity synchrotron map at 23 GHz: The CMB contamination on that map is not likely to correlate so strongly with the best CMB maps available today, after rescaling to 60 GHz. Therefore, the contamination should be in the CMB maps; this would also explain why the synchrotron template is more correlated with the WMAP CMB map than with the TOH one. The relatively high level of contamination might be, potentially, a serious problem for CMB quadrupole C_2^T . Although this point is far from being proved by the present analysis,

the foreground contamination might partly account for the low CMB quadrupole measured by COBE and WMAP.

In spite of the above open problem, we find that the multipole region where there is no evidence for contamination includes the range which dominates the standard WMAP fitting of the reionization optical depth. This result, obtained at 60 GHz, should be adequately representative for the QVW data set. Therefore, it seems really hard to explain in this way the North-South asymmetry in the optical depth fits declared by Hansen et al. (2004a). If such an asymmetry is confirmed, it should be of extragalactic (although not necessarily cosmological) origin. This possibility is supported by Schwartz et al. (2004) in connection with other WMAP anomalies.

Acknowledgments: This work has been carried out in the frame of the SPORt programme funded by the Italian Space Agency (ASI).

REFERENCES

- Bennett, C.L., et al., 2003, ApJS, 148, 97
 Bernardi G., Carretti E., Fabbri R., Sbarra C., Poppi S., Cortiglioni S., 2003, MNRAS, 344, 347
 Bernardi G., Carretti E., Fabbri R., Sbarra C., Poppi S., Cortiglioni S., Jonas J.L., 2004, MNRAS, 351, 436
 Copi C.J., Huterer D., Starkman G.D., 2004, PRD, 70, 043515
 de Oliveira-Costa A., Tegmark M., Zaldarriaga M., Hamilton A., 2004, PRD, 69, 063516
 Dineen P., Coles P., 2004a, MNRAS, 347, 52
 Dineen P., Coles P., 2004b, astro-ph/0410636
 Eriksen H.K. et al., 2004, ApJ, 65, 14
 Hansen F.K., Balbi A., Bandai A.J., Górski K.M., 2004a, MNRAS, 354, 905
 Hansen F.K., Bandai A.J., Górski K.M., 2004b, MNRAS, 354, 665
 Kogut A., et al., 2003, ApJS, 148, 161
 Land K.R., Maguejo J., 2005, astro-ph 0502237
 Naselsky P.D., Doroshkevich A.G., Verkhodanov O.V., 2003, ApJL, 599, 53
 Naselsky P.D., Verkhodanov O.V., Doroshkevich A.G., 2004, MNRAS, 349, 695
 Sbarra C., Carretti E., Cortiglioni S., Zannoni M., Fabbri R., Macculi C., Tucci M., 2003, A&A, 401, 1215
 Schwartz, D.J., Starkman, G.D., Huterer, D., Copi, C.J., 2004, PRL, 93, 221301
 Tegmark M., de Oliveira-Costa A., Hamilton A.J., 2003, PRD, 68, 123523 (TOH)

This paper has been produced using the Royal Astronomical Society/Blackwell Science L^AT_EX style file.